

Multi-GeV Charmonium Mesons

Philip R. Page

Los Alamos National Laboratory

Resonance and Charmonium Studies

Session III, Thu. 3 Aug.

$\bar{p}2000$

Motivation

- Explicit glue tests QCD's non-perturbative gluonic interactions
- Spectroscopy of the new states of matter (gluonic hadrons) essentially unknown
- Charmonium hybrids and glueballs under better theoretical control and experimentally cleaner than light flavour hybrids

Kinematics

p rest

\bar{p} energy E

Centre of mass energy W

$$W^2 = 2m_p (m_p + \sqrt{m_p^2 + E^2})$$

$$E < 10 \text{ GeV} \quad \Rightarrow \quad W < 4500 \text{ MeV}$$

Hybrid Charmonium

Many quenched lattice QCD mass predictions

Decay selection rule:

(non-relativistic quarks, spin 1 pair creation)

P.R. Page, *Phys. Lett.* **B402** (1997) 183

Hybrid charmonium $\xrightarrow{\text{not}} D\bar{D}, D^*\bar{D}^*$
 $\xrightarrow{\text{small}} D^*\bar{D}$

$J^{PC} = 0^{+-} \xrightarrow{\text{not}} D\bar{D}, D^*\bar{D}^*, D^*\bar{D}$

Width $0^{+-}, 1^{-+} = \mathcal{O}(5, 20) \text{ MeV}$


F.E. Close *et al.*, *Phys. Rev.* **D57** (1998) 5653

P.R. Page, *Acta Phys. Polon.* **B29** (1998) 3387

Decay to charmonium + light hadrons signal
 $c\bar{c}$ nature of initial state. ψ easiest to detect

Production: LEAR: $\hat{\rho}(1405), \hat{\rho}(1600)(?), \rho(1450)$

CRYSTAL BARREL



Glueballs

- No light flavour conventional mesons $\gtrsim 3100 \pm 110$ MeV

M.M. Brisudová, L. B., T. G., *Phys. Rev.* **D61** (2000) 054013

- Mixing with charmonium small
- Most glueballs \xrightarrow{not} two glueballs

\implies Glueball could be very narrow!

(if light flavour hybrid meson mixing small)

Production: LEAR: $f_0(1500)$, $f_J(2220)$

Subprogram of hybrid charmonium: Distinguish by charmonium + light hadrons decay

Contents

J^{PC} exotics	J^{PC} unknown
Need PWA	Not need PWA
Fixed \bar{p} energy	Vary \bar{p} energy
Low σ	High σ

J^{PC} Exotics

Conventional charmonium not J^{PC} exotic

J^{PC} exotic \Rightarrow hybrid $c\bar{c}$, glueball (four-quark)

- Need angular distributions / PWA
- + Cannot mix with conventional charmonium

Hybrid $c\bar{c}$: Mass (MeV)

Quenched Lattice QCD:

1^{-+}	0^{+-}	Ref.
$4410^{+60}_{-150} + \text{sys}$	$4560^{+110}_{-100} + \text{sys}$	[1]
$4290^{+110}_{-190} + \text{sys}$	$4560^{+80}_{-110} + \text{sys}$	[1]
$4390 \pm 80 \pm 200$	$4610 \pm 110 \pm 200$	[2]

[1] C. Bernard *et al.* (MILC Collab.), LATTICE'98, hep-lat/9809087

[2] *ibid.*, *Phys. Rev.* **D56** (1997) 7039


$D^{**}D$ threshold: 4290 MeV

2^{+-} , 0^{--} exotics: probably above $D^{**}D$

Hybrid $c\bar{c}$ Production

Formation: $p\bar{p} \xrightarrow{\text{not}} \text{exotic}$

$p\bar{p} \rightarrow \text{exotic} (\pi^0, \pi\pi, \pi\pi\pi, \eta, \dots)$



\Rightarrow Fixed \bar{p} energy

$$E < 10 \text{ GeV} \quad \Rightarrow \quad m_{\text{exotic}} < 4360 \text{ MeV}$$

Hybrid $c\bar{c}$: Decay

$$1^{-+} \rightarrow \psi (\omega, \phi, \gamma) \rightarrow e^+ e^- e^+ e^-, e^+ e^- \gamma$$

[small]

$$0^{+-} \rightarrow \psi (\pi\pi)_S [\psi\pi^0\pi^0] \rightarrow e^+ e^- \gamma\gamma\gamma\gamma$$

[background]

Missing mass:

Detect π^0 and ψ in $p\bar{p} \rightarrow$ exotic $\pi^0 \rightarrow \psi X \pi^0$

J^{PC}	Open charm	Hidden charm	Light hadrons
0^{+-}	Quantum numbers forbid $D^{(*)}D^{(*)}$	$J/\psi\{f_{\{0,1,2\}}, (\pi\pi)_S\}$ $h_c\eta; \eta_c h_1$ $\chi_{c0}\omega$ $\chi_{c\{1,2\}}\{\omega, h_1, \gamma\}$	$a_{\{0,1,2\}}\rho; a_{\{1,2\}}\{b_1, \gamma\}$ $b_1\pi; h_1\eta^{(')}$ $\{(\pi\pi)_S, f_0\}\{\omega, \phi\}$ $f_{\{1,2\}}\{\omega, h_1, \phi, \gamma\}$
0^{--}	D^*D	$h_c(\pi\pi)_S$ $J/\psi\{f_{\{1,2\}}, \eta^{(')}\}$ $\chi_{c0}h_1; \eta_c\{\omega, \phi\}$ $\chi_{c\{1,2\}}\{\omega, h_1, \gamma\}$	$a_{\{0,1,2\}}b_1; a_{\{1,2\}}\{\rho, \gamma\}$ $\rho\pi$ $f_0h_1; \eta^{(')}\{\omega, \phi\}$ $f_{\{1,2\}}\{\omega, h_1, \phi, \gamma\}$
1^{-+}	D^*D, D^*D^*	$\chi_{c\{0,1,2\}}(\pi\pi)_S$ $\eta_c\{f_{\{1,2\}}, \eta^{(')}\}$ $\chi_{c\{1,2\}}\eta$ $\{h_c, J/\psi\}\{\omega, h_1, \phi, \gamma\}$	$a_{\{0,1,2\}}a_{\{0,1,2\}}; a_{\{1,2\}}\pi$ $f_{\{0,1,2\}}f_{\{0,1,2\}}; f_{\{1,2\}}\eta^{(')}$ $\{\rho, \gamma\}\{\rho, b_1\}; b_1b_1$ $\{\omega, h_1, \phi, \gamma\}\{\omega, h_1, \phi, \gamma\}$
2^{+-}	D^*D, D^*D^*	$\{h_c, J/\psi\}\{f_{\{0,1,2\}}, (\pi\pi)_S\}$ $\{h_c, J/\psi\}\eta^{(')}$ $\{\eta_c, \chi_{c\{0,1,2\}}\}\{\omega, h_1, \phi, \gamma\}$	$a_{\{0,1,2\}}\{\rho, b_1, \gamma\}$ $\{\rho, \gamma, b_1\}\pi$ $\{\eta^{(')}, f_{\{0,1,2\}}\}\{\omega, h_1, \phi, \gamma\}$

Hybrid $c\bar{c}$: Cross-section

LEAR: similar to charmonium: 130 pb


$\sigma(p\bar{p} \rightarrow \psi\pi^0) = 130 \pm 25$ pb. from R. Cester, Workshop on Physics
at SuperLEAR (Zurich, Oct. 1991), Inst. of Physics Conf. Series
No. 124, p. 91

$$\mathcal{L} = 10^{32} \text{ cm}^{-2} \text{ s}^{-1} \quad 50\% \text{ efficiency}$$

$$BR(\text{exotic} \rightarrow \psi\omega) = 1\%$$

\Rightarrow 1 event/day

Glueballs



- + Large σ
- Many decay channels: light hadrons

$2^{+-} \quad 4140 \pm 50 \pm 200 \text{ MeV}$ [lightest]

$0^{+-} \quad 4740 \pm 70 \pm 230 \text{ MeV}$ [too heavy]

C.J. Morningstar, M. Peardon, *Phys. Rev.* **D60** (1999) 034509

J^{PC} Unknown

- + PWA not necessary \Rightarrow bump hunting
- Can be conventional charmonium
- + Narrowness or decay modes distinctive
- + Small conventional charmonium mixing, except for mass coincidences $[1^{--}]$

S.B. Gerasimov, *Proc. of 11th Int. Conf. on Problems of Quantum Field Theory* (July 1998, Dubna, Russia), hep-ph/9812509

F.E. Close, P.R. Page, *Phys. Lett.* **B366** (1996) 323

Hybrid $c\bar{c}$: Mass

8 low-lying hybrids [5 non-exotic]:

$$1^{--}, (0, 1, 2)^{-+} < 1^{++}, (0, 1, 2)^{+-}$$

K.J. Juge, J. Kuti, C.J. Morningstar, hep-lat/9909165

T. Manke, *et al.*, *Phys. Rev.* **D57** (1998) 3829

Splittings (MeV)

J^{PC}	Gluon exchange [1]	Confinement [2]
0^{-+}	-180	8
1^{-+}	-50	4
1^{--}	60	0
2^{-+}	210	-4

[1] P. R. Page, *D. Phil. thesis* (Univ. of Oxford, 1995)

[2] J. Merlin, J. Paton, *Phys. Rev.* **D35** (1987) 1668

Hybrid $c\bar{c}$: Production

Formation: $p\bar{p} \rightarrow$ non-exotic $W < 4500$ MeV
⇒ Energy scan

Hybrid $c\bar{c}$: Decay

$p\bar{p} \rightarrow$ non-exotic $\rightarrow \psi X$. ψ detected. X from missing mass. $X = \eta, \eta', \omega, \phi, \pi\pi, K\bar{K}, \dots$
[conventional charmonium suppressed]

$\psi\eta$ $1^{--}, 1^{+-}$

$\psi\omega$ $0^{-+}, 2^{-+}, 1^{++}$

non-exotic $\rightarrow e^+e^-, \mu^+\mu^-, \tau^+\tau^-, \gamma\gamma$ small

S. Ono *et al.*, *Z. Phys. C26* (1984) 307; *Phys. Rev. D34* (1986) 186; P.R. Page, *Nucl. Phys. B495* (1997) 268

Hybrid $c\bar{c}$: Cross-section

Form. $\sigma = 0.1 \mu\text{b}$. Similar to charmonium

F.E. Close, SuperLEAR Workshop, p. 63; K. Königsmann, p. 71

$$\mathcal{L} = 10^{32} \text{ cm}^{-2} \text{ s}^{-1} \quad 50\% \text{ efficiency}$$


$$BR(\text{exotic} \rightarrow \psi\eta) = 1\%$$

\Rightarrow 500 events/day

No PWA \Rightarrow can detect all non-exotic $\rightarrow \psi X$

$\approx 25\% \Rightarrow 12500$ events/day

Glueballs



- + Large σ
- Many decay channels: light hadrons

$1^{--} \quad 3850 \pm 50 \pm 190$ MeV

$2^{-+} \quad 3890 \pm 40 \pm 190$ MeV


$2^{--} \quad 3930 \pm 40 \pm 190$ MeV

$3^{--} \quad 4130 \pm 90 \pm 200$ MeV

C.J. Morningstar, M. Peardon, *Phys. Rev.* **D60** (1999) 034509

non-exotic $\rightarrow e^+e^-$, $\mu^+\mu^-$, $\tau^+\tau^-$, $\gamma\gamma$ small

Bump Hunting



M.M. Brisudová, L. B., T. G., *Phys. Rev.* **D61** (2000) 054013
K. Heikkilä, N.A. Törnqvist, S. Ono, *Phys. Rev.* **D29** (1984) 110

18 conventional charmonia \Rightarrow

contiguous resonances if $c\bar{c}$ width ≈ 50 MeV

$\gtrsim 4$ non-exotic glueballs

$\lesssim 5$ non-exotic hybrid charmonia

Given decay channel will be J^{PC} filter

Example: $\psi\eta$: 3 charmonia [S-, P-wave]

2 hybrid charmonia

2 glueballs [not preferred]

Need good understanding of conventional states
so that new states stand out

Energy Scan 4.0 - 4.3 GeV region:

10 MeV bins $\Leftrightarrow \bar{p}$ tuned 50 MeV \Rightarrow 30 scans

30 MeV bins $\Leftrightarrow \bar{p}$ tuned 150 MeV \Rightarrow 10 scans

Lowest mass new flavoured baryons:
Doubly charmed

$$W \gtrsim 2 \times 3650 \text{ MeV} \Rightarrow E \gtrsim 27 \text{ GeV}$$

Lowest mass new flavoured mesons:
Bottom charm

$$W \geq 2 \times (6.40 \pm 0.39 \pm 0.13) \text{ GeV} \Rightarrow E \gtrsim 86 \text{ GeV}$$

D.E. Groom *et al.*, *Eur. Phys. J. C15* (2000) 1

Competition

Proposed Glue / Charm - Factory @ GSI Darmstadt: $E < 15$ GeV

Conclusions

J^{PC} exotics	J^{PC} unknown
Need PWA	Not need PWA
Fixed \bar{p} energy	Vary \bar{p} energy
Low σ	High σ